

Highlights on n-rich Λ -Hypernuclei from the FINUDA experiment

E. Botta

INFN - Sezione di Torino and Torino University - Physics Department
via P. Giuria 1, Torino, Italy

Λ -hypernuclei are long lived systems in which the Λ hyperon acts as constituent nucleon. The strangeness degree of freedom of the hyperon makes the Pauli exclusion principle ineffective on it and allows the Λ to populate sharp single particle shell model states down to the $\Lambda(1s)$ ground state.

In Λ -hypernuclei, then, a strong contribution to the total binding energy is added by the Λ , with the possibility of producing bound systems containing unstable core nuclei (glue-like rôle of the Λ). Λ -hypernuclei are thus suitable tools for investigating neutron-rich (proton-rich) systems, even beyond the neutron (proton) drip line.

The study of neutron-rich Λ -hypernuclei is one the main topics of the scientific program of the FINUDA experiment which has completed its operation at DAΦNE, the INFN-LNF (e^+ , e^-) collider working at the $\Phi(1020)$ center of mass energy. FINUDA has performed an extensive search for bound neutron-rich Λ -hypernuclear states. In the first data taking (2003-2004), ${}^6_\Lambda\text{H}$, ${}^7_\Lambda\text{H}$ and ${}^{12}_\Lambda\text{Be}$ have been investigated by looking at the π^+ from the double charge exchange (K_{stop}^-, π^+) production reaction; upper limits for the production rates have been reported [1]: $R_{\pi^+}({}^6_\Lambda\text{H}) < (2.5 \pm 0.4_{\text{stat}}^{+0.4}_{-0.1\text{syst}}) \cdot 10^{-5}/K_{stop}^-$, $R_{\pi^+}({}^7_\Lambda\text{H}) < (4.5 \pm 0.9_{\text{stat}}^{+0.4}_{-0.1\text{syst}}) \cdot 10^{-5}/K_{stop}^-$ and $R_{\pi^+}({}^{12}_\Lambda\text{Be}) < (2.0 \pm 0.4_{\text{stat}}^{+0.3}_{-0.1\text{syst}}) \cdot 10^{-5}/K_{stop}^-$.

In the second data taking (2006-2007) a ~ 5 times larger statistics has been collected and a new analysis technique has been applied for the search of bound ${}^6_\Lambda\text{H}$ and ${}^9_\Lambda\text{He}$, based on the coincidence between a π^+ from the (K_{stop}^-, π^+) production reaction and a π^- from the two body mesonic weak decay of the hypernucleus to $\pi^- + {}^6\text{He}_{g.s.}$. With this method the existence of ${}^6_\Lambda\text{H}$ as a bound state has been assessed, based on 3 clearly identified events, with a binding energy $B_\Lambda = (4.0 \pm 1.1)$ MeV, with respect to ${}^5\text{H} + \Lambda$; and a production rate $R_{\pi^+}({}^6_\Lambda\text{H}) = (5.9 \pm 4.0) \cdot 10^{-6}/K_{stop}^-$ [2,3] has been evaluated. Indications on the structure of the ${}^6_\Lambda\text{H}$ energy levels have been obtained from a systematic difference among the ${}^6_\Lambda\text{H}$ mass evaluated from production and decay reactions.

The same method has been applied to the search for bound ${}^9_\Lambda\text{He}$, which is interesting since it could be a neutron-halo Hypernucleus. No event was found and an upper limit for its production, $R_{\pi^+}({}^9_\Lambda\text{He}) < (5.0 \pm 4.1) \cdot 10^{-6}/K_{stop}^-$, was deduced [4].

In the presentation a review of all observed neutron-rich Λ -hypernuclei will be given, with particular emphasis on the FINUDA results and on their discussion.

[1] M. Agnello, et al., Physics Letters B 640 (2006) 145.

[2] M. Agnello et al., Physical Review Letters 108 (2012) 042501.

[3] M. Agnello et al., Nuclear Physics A 881 (2012) 269.

[4] M. Agnello et al., Physical Review C 86 (2012) 057301.

E-mail:

botta@to.infn.it